



## Guidance Document for the Prehospital Use of Tranexamic Acid in Injured Patients

Peter E. Fischer MD, MS, FACS, Eileen M. Bulger MD, FACS, Debra G. Perina MD, FACEP, Theodore R. Delbridge MD, MPH, Mark L. Gestring MD, FACS, Mary E. Fallat MD, FACS, David V. Shatz MD, FACS, Jay Doucet MD, MSc, FACS, Michael Levy MD, FACEP, Lance Stuke MD, MPH, FACS, Scott P. Zietlow MD, FACS, Jeffrey M. Goodloe MD, NRP, Wayne E. VanderKolk MD, Adam D. Fox DPM, DO, FACS & Nels D. Sanddal PhD, NREMT

To cite this article: Peter E. Fischer MD, MS, FACS, Eileen M. Bulger MD, FACS, Debra G. Perina MD, FACEP, Theodore R. Delbridge MD, MPH, Mark L. Gestring MD, FACS, Mary E. Fallat MD, FACS, David V. Shatz MD, FACS, Jay Doucet MD, MSc, FACS, Michael Levy MD, FACEP, Lance Stuke MD, MPH, FACS, Scott P. Zietlow MD, FACS, Jeffrey M. Goodloe MD, NRP, Wayne E. VanderKolk MD, Adam D. Fox DPM, DO, FACS & Nels D. Sanddal PhD, NREMT (2016) Guidance Document for the Prehospital Use of Tranexamic Acid in Injured Patients, *Prehospital Emergency Care*, 20:5, 557-559, DOI: [10.3109/10903127.2016.1142628](https://doi.org/10.3109/10903127.2016.1142628)

To link to this article: <http://dx.doi.org/10.3109/10903127.2016.1142628>



Published online: 17 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 665



View related articles [↗](#)



View Crossmark data [↗](#)

# SPECIAL CONTRIBUTION

## GUIDANCE DOCUMENT FOR THE PREHOSPITAL USE OF TRANEXAMIC ACID IN INJURED PATIENTS

Peter E. Fischer, MD, MS, FACS, Eileen M. Bulger, MD, FACS, Debra G. Perina, MD, FACEP, Theodore R. Delbridge, MD, MPH, Mark L. Gestring, MD, FACS, Mary E. Fallat, MD, FACS, David V. Shatz, MD, FACS, Jay Doucet, MD, MSc, FACS, Michael Levy, MD, FACEP, Lance Stuke, MD, MPH, FACS, Scott P. Zietlow, MD, FACS, Jeffrey M. Goodloe, MD, NRP, FACEP, Wayne E. VanderKolk, MD, Adam D. Fox, DPM, DO, FACS, Nels D. Sanddal, PhD, NREMT

---

Received December 31, 2015 from Division of Trauma/Surgical Critical Care and Acute Care Surgery, Department of Surgery, Carolinas Medical Center, Charlotte, North Carolina (PEF); Division of Trauma, Department of Surgery, University of Washington, Harborview Medical Center, Seattle, Washington (EMB); Department of Emergency Medicine, University of Virginia, Charlottesville, Virginia (DGP); Department of Emergency Medicine, East Carolina University, Greenville, North Carolina (TRD); Department of Surgery, University of Pittsburgh Medical Center Health System, Pittsburgh, Pennsylvania (MG); Division of Pediatric Surgery, Department of Surgery, University of Louisville Hospital, Louisville, Kentucky (MEF); Division of Trauma, Acute Care Surgery and Surgical Critical Care, Department of Surgery, University of California Davis Medical Center, Sacramento, California (DVS); Department of Surgery, University of California San Diego Health System, San Diego, California (JD); Anchorage Fire Department, Anchorage, Alaska (ML); Department of Surgery, Louisiana State University, New Orleans, Louisiana (LS); Division of Trauma, Critical Care and General Surgery, Department of Surgery, Mayo Clinic, Rochester, Minnesota (SPZ); Department of Emergency Medicine, The University of Oklahoma School of Community Medicine, Tulsa, Oklahoma (JMG); St. Mary Trauma Service, Grand Rapids, Michigan (WEV); Department of Surgery, Rutgers New Jersey Medical School, Newark, New Jersey (ADF); American College of Surgeons, Chicago, Illinois (NDS). Accepted for publication January 12, 2016.

Endorsed by the American College of Surgeons–Committee on Trauma; American College of Emergency Physicians, and National Association of EMS Physicians

The authors have no conflicts of interest or financial support to disclose.

Address correspondence to Peter E. Fischer, MD, MS, Medical Director, Metrolina Trauma Advisory Committee, Trauma/Surgical Critical Care & Acute Care Surgery, F.H. “Sammy” Ross Jr. Trauma Center, Carolinas Medical Center, P.O. Box 32861, Charlotte, NC 28232-2861, USA. E-mail: Peter.Fischer@carolinashealthcare.org

doi: 10.3109/10903127.2016.1142628

### ABSTRACT

Tranexamic acid (TXA) is being administered already in many prehospital air and ground systems. Insufficient evidence exists to support or refute the prehospital administration of TXA, and results are pending from several prehospital studies currently in progress. We have created this document to aid agencies and systems in best practices for TXA administration based on currently available best evidence. This document has been endorsed by the American College of Surgeons–Committee on Trauma, the American College of Emergency Physicians, and the National Association of EMS Physicians. **Key words** tranexamic acid; trauma; hemorrhage control; emergency medical services

PREHOSPITAL EMERGENCY CARE 2016;20:557–559

### INTRODUCTION

Tranexamic acid (TXA) is a synthetic lysine analog that competitively inhibits the activation of plasminogen to plasmin. It was approved by the U.S. Food and Drug Administration in 1986 for short-term use as an injection to reduce or prevent bleeding during tooth extraction in hemophilia patients. The oral form is approved to treat menorrhagia. All other uses are off-label.

TXA has the potential benefit of decreasing fibrinolysis in the injured, bleeding patient. The 2010 results of the Clinical Randomization of an Antifibrinolytic in Significant Hemorrhage (CRASH-2) trial showed a decrease in all-cause mortality of 1.5% and a decrease in risk of death secondary to hemorrhage of 0.8% in patients receiving TXA.<sup>1</sup> While CRASH-2 was the largest randomized study to date, many are critical of the findings, such as the lack of modern resuscitative practices in many of the participating centers.<sup>2</sup> Retrospective studies, such as the Military Application of Tranexamic Acid in Trauma Emergency Resuscitation (MATTERS)

study,<sup>3</sup> also showed a mortality benefit in patients receiving TXA but suggested an increase in thromboembolic events.

A subgroup analysis of the CRASH-2 study examined the timing of administration.<sup>4</sup> In bleeding patients, TXA administration within 1 hour of injury resulted in a significant reduction in mortality. This benefit continued up to 3 hours post-injury; however, after 3 hours, TXA administration was found to be harmful. Some in the prehospital community have embraced this finding, suggesting that an early administration approach is most beneficial to the patient and, thus, the drug is ideal for the prehospital environment. Our organizations recognize that TXA is being administered already in many prehospital air and ground systems. We recognize that insufficient evidence exists to support or refute the prehospital administration of TXA and are awaiting the results of several prehospital studies currently in progress before making a recommendation regarding prehospital use of this agent. Therefore, we have created this document to aid agencies and systems in best practices for TXA administration based on currently available best evidence.

### SYSTEM INTEGRATION IS PARAMOUNT

Prompt hemorrhage control and rapid transport to a trauma center are both key to survival in the bleeding patient. TXA administration should never delay transport. Furthermore, TXA administration involves a bolus dose followed by an infusion over 8 hours. Prior to the implementation of field TXA, the EMS agency and the receiving trauma center should jointly develop protocols to ensure eligible patients receive the appropriate bolus dose in the field and infusion dose at the hospital. A clear hand-off care report given by EMS providers is paramount, with specific notation that the TXA bolus has been given during transport to avoid repeat bolus doses and to ensure that infusion is initiated in a timely fashion.

### ADMINISTRATION TO BLEEDING PATIENTS

Limited evidence suggests that more venous thromboembolic events (VTE) occur when TXA is given to patients not requiring massive transfusion.<sup>5</sup> This discrimination can be difficult in the field where rapid decisions must be made with minimal diagnostics or time for evaluation. Objective measurements should be used to guide prehospital TXA administration protocols. The focus for management of compressible, external bleeding should be on direct pressure, tourniquets, hemostatic agents, and/or wound packing. Evidence of injury consistent with non-compressible hemorrhage (e.g., penetrating thoracoabdominal trauma or unstable pelvis fractures) along with heart rate >120

bpm and systolic blood pressure (SBP) <90 mmHg are suggested criteria. Agencies may consider vital sign adjustments for the geriatric population.

### TRANSPORT TO A TRAUMA CENTER

Compressible bleeding should be managed with pressure dressings, hemostatic agents, wound packing, or tourniquets. Prehospital TXA should be administered only to patients with non-compressible bleeding. Ultimately, definitive surgical control at a trauma center is essential. Patients receiving prehospital TXA should be preferentially transported to a Level I or II trauma center if available. If geographic or other factors preclude direct trauma center transport, the first receiving hospital should be capable of continuing the TXA infusion and implementing hemorrhage-control procedures.

### MONITORING AND QUALITY IMPROVEMENT

Given the lack of data available, our organizations recommend that prehospital TXA administration be monitored closely in a prehospital and/or trauma registry. Administration should be reviewed and protocols constantly refined to avoid unnecessary or incomplete doses, inappropriate patient selection, or lack of infusion following the initial bolus. TXA dosing, timing, blood transfusion requirements, and any adverse events should be included in the registry. If thromboelastography (TEG) is used to guide resuscitation, fibrinolysis as measured by percent clot lysis at 30 minutes after reaching maximum clot strength (LY30) should be collected in the trauma registry for future research.

### CAUTION IN CASES OF KNOWN ANTICOAGULATION

The effect of administering TXA in conjunction with other medications used to reverse anticoagulants (e.g. FEIBA, an anti-inhibitor coagulant complex) may lead to more thrombotic complications. In the case of trauma patients known to be on chronic anticoagulants, medical control should be consulted prior to TXA administration if specific protocols for this situation have not been developed.

### COMMUNICATION IS KEY

Repeat bolus doses of TXA should be avoided. Prehospital TXA administration should be clearly communicated with the next receiving provider. Simple adjuncts, such as stickers or wristbands applied to patients, may be used to aid in the information transfer.

## PEDIATRICS

Children were excluded from the CRASH-2 Trial.<sup>1</sup> The best studies of TXA efficacy in children to date are in craniofacial surgery, where there is Level I evidence that TXA decreases transfusion requirements.<sup>6–8</sup> An acknowledged risk of TXA is seizures, which has been seen mainly in the pediatric cardiac surgery population. With the extremely limited information concerning TXA use in pediatric trauma, prehospital TXA use in pediatric patients is not currently recommended outside of a research study.

## DON'T FORGET THE BASICS

In the bleeding patient, hemorrhage control and appropriate resuscitation remain the priority. Prehospital TXA use should never supersede field bleeding control techniques, rapid transport to a trauma center, or the administration of blood or plasma.

## References

1. Crash- Trial Collaborators, Shakur H, Roberts I, et al. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet*. 2010;376(9734):23–32.
2. Napolitano LM, Cohen MJ, Cotton BA, Schreiber MA, Moore EE. Tranexamic acid in trauma: how should we use it? *J Trauma Acute Care Surg*. 2013;74(6):1575–86.
3. Morrison JJ, Dubose JJ, Rasmussen TE, Midwinter MJ. Military Application of Tranexamic Acid in Trauma Emergency Resuscitation (MATTERs) Study. *Arch Surg*. 2012;147(2):113–19.
4. Crash- Trial Collaborators, Roberts I, Shakur H, et al. The importance of early treatment with tranexamic acid in bleeding trauma patients: an exploratory analysis of the CRASH-2 randomised controlled trial. *Lancet*. 2011;377(9771):1096–101, 1101 e1091–92.
5. Stockinger Z, Gross KR, Butler FK, Bailey JA. Does TXA Matter? Refining Indications For its Use in Combat Casualties. Paper presented at: Eastern Association for the Surgery of Trauma Annual Scientific Assembly; January 13–17, 2015; Lake Buena Vista, Florida.
6. Goobie SM, Meier PM, Pereira LM, et al. Efficacy of tranexamic acid in pediatric craniostylosis surgery: a double-blind, placebo-controlled trial. *Anesthesiology*. 2011;114(4):862–71.
7. Dadure C, Sauter M, Bringuier S, et al. Intraoperative tranexamic acid reduces blood transfusion in children undergoing craniostylosis surgery: a randomized double-blind study. *Anesthesiology*. 2011;114(4):856–61.
8. Faraoni D, Goobie SM. The efficacy of antifibrinolytic drugs in children undergoing noncardiac surgery: a systematic review of the literature. *Anesth Analg*. 2014;118(3):628–36.